

## AGRICULTURE

# A global synthesis reveals biodiversity-mediated benefits for crop production

Matteo Dainese<sup>1,2\*</sup>, Emily A. Martin<sup>2</sup>, Marcelo A. Aizen<sup>3</sup>, Matthias Albrecht<sup>4</sup>, Ignasi Bartomeus<sup>5</sup>, Riccardo Bommarco<sup>6</sup>, Luisa G. Carvalheiro<sup>7,8</sup>, Rebecca Chaplin-Kramer<sup>9</sup>, Vesna Gagic<sup>10</sup>, Lucas A. Garibaldi<sup>11</sup>, Jaboury Ghazoul<sup>12</sup>, Heather Grab<sup>13</sup>, Mattias Jonsson<sup>6</sup>, Daniel S. Karp<sup>14</sup>, Christina M. Kennedy<sup>15</sup>, David Kleijn<sup>16</sup>, Claire Kremen<sup>17</sup>, Douglas A. Landis<sup>18</sup>, Deborah K. Letourneau<sup>19</sup>, Lorenzo Marini<sup>20</sup>, Katja Poveda<sup>13</sup>, Romina Rader<sup>21</sup>, Henrik G. Smith<sup>22,23</sup>, Teja Tschardt<sup>24</sup>, Georg K. S. Andersson<sup>22</sup>, Isabelle Badenhausser<sup>25,26</sup>, Svenja Baensch<sup>24,27</sup>, Antonio Diego M. Bezerra<sup>28</sup>, Felix J. J. A. Bianchi<sup>29</sup>, Virginie Boreux<sup>12,30</sup>, Vincent Bretagnolle<sup>31</sup>, Berta Caballero-Lopez<sup>32</sup>, Pablo Cavigliasso<sup>33</sup>, Aleksandar Ćetković<sup>34</sup>, Natacha P. Chacoff<sup>35</sup>, Alice Classen<sup>2</sup>, Sarah Cusser<sup>36</sup>, Felipe D. da Silva e Silva<sup>37,38</sup>, G. Arjen de Groot<sup>39</sup>, Jan H. Dudenhöffer<sup>40</sup>, Johan Ekroos<sup>22</sup>, Thijs Fijen<sup>16</sup>, Pierre Franck<sup>41</sup>, Breno M. Freitas<sup>28</sup>, Michael P. D. Garratt<sup>42</sup>, Claudio Gratton<sup>43</sup>, Juliana Hipólito<sup>11,44</sup>, Andrea Holzschuh<sup>2</sup>, Lauren Hunt<sup>45</sup>, Aaron L. Iverson<sup>13</sup>, Shalene Jha<sup>46</sup>, Tamar Keasar<sup>47</sup>, Tania N. Kim<sup>48</sup>, Miriam Kishinevsky<sup>49</sup>, Björn K. Klatt<sup>23,24</sup>, Alexandra-Maria Klein<sup>30</sup>, Kristin M. Krewenka<sup>50</sup>, Smitha Krishnan<sup>12,51,52</sup>, Ashley E. Larsen<sup>53</sup>, Claire Lavigne<sup>41</sup>, Heidi Lieke<sup>54</sup>, Bea Maas<sup>55</sup>, Rachel E. Mallinger<sup>56</sup>, Eliana Martinez Pachon<sup>57</sup>, Alejandra Martínez-Salinas<sup>58</sup>, Timothy D. Meehan<sup>59</sup>, Matthew G. E. Mitchell<sup>60</sup>, Gonzalo A. R. Molina<sup>61</sup>, Maike Nesper<sup>12</sup>, Lovisa Nilsson<sup>22</sup>, Megan E. O'Rourke<sup>62</sup>, Marcell K. Peters<sup>2</sup>, Milan Plečáček<sup>34</sup>, Simon G. Potts<sup>43</sup>, Davi de L. Ramos<sup>63</sup>, Jay A. Rosenheim<sup>64</sup>, Maj Rundlöf<sup>23</sup>, Adrien Rusch<sup>65</sup>, Agustín Sáez<sup>66</sup>, Jeroen Scheper<sup>16,39</sup>, Matthias Schleuning<sup>67</sup>, Julia M. Schmack<sup>68</sup>, Amber R. Sciligo<sup>69</sup>, Colleen Seymour<sup>70</sup>, Dara A. Stanley<sup>71</sup>, Rebecca Stewart<sup>22</sup>, Jane C. Stout<sup>72</sup>, Louis Sutter<sup>4</sup>, Mayura B. Takada<sup>73</sup>, Hisatomo Taki<sup>74</sup>, Giovanni Tamburini<sup>30</sup>, Matthias Tschumi<sup>4</sup>, Blandina F. Viana<sup>75</sup>, Catrin Westphal<sup>27</sup>, Bryony K. Willcox<sup>21</sup>, Stephen D. Wratten<sup>76</sup>, Akira Yoshioka<sup>77</sup>, Carlos Zaragoza-Trello<sup>5</sup>, Wei Zhang<sup>78</sup>, Yi Zou<sup>79</sup>, Ingolf Steffan-Dewenter<sup>2</sup>

Human land use threatens global biodiversity and compromises multiple ecosystem functions critical to food production. Whether crop yield-related ecosystem services can be maintained by a few dominant species or rely on high richness remains unclear. Using a global database from 89 studies (with 1475 locations), we partition the relative importance of species richness, abundance, and dominance for pollination; biological pest control; and final yields in the context of ongoing land-use change. Pollinator and enemy richness directly supported ecosystem services in addition to and independent of abundance and dominance. Up to 50% of the negative effects of landscape simplification on ecosystem services was due to richness losses of service-providing organisms, with negative consequences for crop yields. Maintaining the biodiversity of ecosystem service providers is therefore vital to sustain the flow of key agroecosystem benefits to society.

## INTRODUCTION

Natural and modified ecosystems contribute a multitude of functions and services that support human well-being (1, 2). It has long been recognized that biodiversity plays an important role in the functioning of ecosystems (3, 4), but the dependence of ecosystem services on biodiversity is under debate. An early synthesis revealed inconsistent results (5), whereas subsequent studies suggest that a few dominant species may supply the majority of ecosystem services (6, 7). It thus remains unclear whether a few dominant or many complementary species are needed to supply ecosystem services.

The interpretation of earlier studies has been controversial because multiple mechanisms underlying changes in ecosystem service response to biodiversity can operate in combination (8, 9). On one hand, communities with many species are likely to include species responsible for large community-wide effects due to statistical selection. On the other hand, such diverse communities may contain a particular combination

of species that complement each other in service provisioning. While these mechanisms imply positive effects of species richness on ecosystem service supply, total organism abundance or dominance of certain species may also drive the number of interactions benefiting ecosystem service supply. Depending on the relative importance of species complementarity, community abundance, and the role of dominant species, different relationships between species richness and ecosystem services can be expected (10).

In real-world ecosystems, natural communities consist of a few highly abundant (dominant species) and many rare ones. The importance of richness, abundance, and dominance is likely to be influenced by the extent to which relative abundance changes with species richness (11) and by differences in the effectiveness and degree of specialization of service-providing communities. However, these three aspects of diversity have typically been tested in isolation and mainly in small-scale experimental settings (12, 13), while a synthetic study contrasting their

Copyright © 2019  
The Authors, some  
rights reserved;  
exclusive licensee  
American Association  
for the Advancement  
of Science. No claim to  
original U.S. Government  
Works. Distributed  
under a Creative  
Commons Attribution  
NonCommercial  
License 4.0 (CC BY-NC).

Downloaded from <http://advances.sciencemag.org/> on October 16, 2019

relative importance in real-world ecosystems is still lacking. A major limitation to resolving these relationships is a lack of evidence from real-world human-driven biodiversity changes (14, 15), particularly for ecosystem services in agroecosystems. For instance, changes in richness and total or relative abundance of service-providing organisms in response to land-clearing for agriculture (16, 17) could alter the flow of benefits to people in different ways compared to experimental random loss of biodiversity.


Over the past half-century, the need to feed a growing world population has led to markedly expanded and intensified agricultural production, transforming many regions into simplified landscapes (18). This transformation not only has contributed to enhanced agricultural production but also has led to the degradation of the global environment. The loss of biodiversity can disrupt key intermediate services to agriculture, such as crop pollination (19) and biological pest control (20), which underpin the final provisioning service of crop

production (21). The recent stagnation or even decline of crop yields with ongoing intensification (22) indicates that alternative pathways are necessary to maintain future stable and sustainable crop production (23–25). An improved understanding of global biodiversity-driven ecosystem services in agroecosystems and their cascading effects on crop production is urgently needed to forecast future supplies of ecosystem services and to pursue strategies for sustainable management (15).

We compiled an extensive database comprising 89 studies that measured richness and abundance of pollinators, pest natural enemies, and associated ecosystem services at 1475 sampling locations around the world (Fig. 1A). We focused on the ecosystem services of pollination and biological pest control because these services are essential to crop production and have been the focus of much research in recent decades (26). We quantified pollinator and pest natural enemy richness as the number of unique taxa sampled from each location (field), abundance as the number of observed individuals, and evenness (or the complementary term,

<sup>1</sup>Institute for Alpine Environment, Eurac Research, Viale Druso 1, 39100 Bozen/Bolzano, Italy. <sup>2</sup>Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Am Hubland, 97074 Würzburg, Germany. <sup>3</sup>Grupo de Ecología de la Polinización, INIBIOMA, Universidad Nacional del Comahue, CONICET, 8400 Bariloche, Río Negro, Argentina. <sup>4</sup>Agroecology and Environment, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland. <sup>5</sup>Estación Biológica de Doñana (EBD-CSIC), Integrative Ecology, E-41092 Sevilla, Spain. <sup>6</sup>Swedish University of Agricultural Sciences, Department of Ecology, 750 07 Uppsala, Sweden. <sup>7</sup>Departamento de Ecología, Universidade Federal de Goiás (UFG), Goiânia, Brazil. <sup>8</sup>Faculdade de Ciências, Centre for Ecology, Evolution and Environmental Changes (CE3C), Universidade de Lisboa, Lisboa, Portugal. <sup>9</sup>Natural Capital Project, Stanford University, Stanford, CA 94618, USA. <sup>10</sup>CSIRO, GPO Box 2583, Brisbane, QLD 4001, Australia. <sup>11</sup>Instituto de Investigaciones en Recursos Naturales, Agroecología y Desarrollo Rural (INRAD), Sede Andina, Universidad Nacional de Río Negro (UNRN) y CONICET, Mitre 630, CP 8400 San Carlos de Bariloche, Río Negro, Argentina. <sup>12</sup>Department of Environmental Systems Science, ETH Zurich, Universitätstrasse 16, 8092 Zurich, Switzerland. <sup>13</sup>Department of Entomology, Cornell University, Ithaca, NY 14853, USA. <sup>14</sup>Department of Wildlife, Fish and Conservation Biology, University of California Davis, Davis, CA 95616, USA. <sup>15</sup>Global Lands Program, The Nature Conservancy, 117 E. Mountain Avenue, Fort Collins, CO 80524, USA. <sup>16</sup>Plant Ecology and Nature Conservation Group, Wageningen University, Droevendaalsesteeg 3a, Wageningen 6708 PB, Netherlands. <sup>17</sup>IREC and Biodiversity Research Centre, University of British Columbia, Vancouver, BC, Canada. <sup>18</sup>Department of Entomology and Great Lakes Bioenergy Research Center, Michigan State University, 204 CIPS, 578 Wilson Ave, East Lansing, MI 48824, USA. <sup>19</sup>Department of Environmental Studies, University of California, Santa Cruz, Santa Cruz, CA 95064, USA. <sup>20</sup>DAFNAE, University of Padova, viale dell'Università 16, 35020 Legnaro, Padova, Italy. <sup>21</sup>School of Environment and Rural Science, University of New England, Armidale, NSW 2350, Australia. <sup>22</sup>Centre for Environmental and Climate Research, Lund University, S-223 62 Lund, Sweden. <sup>23</sup>Department of Biology, Lund University, S-223 62 Lund, Sweden. <sup>24</sup>Agroecology, Department of Crop Sciences, University of Göttingen, D-37077 Göttingen, Germany. <sup>25</sup>USC1339 INRA-CNRS, CEBC UMR 7372, CNRS and Université de La Rochelle, Beauvoir sur Niort 79360, France. <sup>26</sup>INRA, Unité de Recherche Pluridisciplinaire Prairies et Plantes Fourragères (URP3F), Lusignan 86600, France. <sup>27</sup>Functional Agrobiodiversity, Department of Crop Sciences, University of Göttingen, Germany. <sup>28</sup>Departamento de Zootecnia—CCA, Universidade Federal do Ceará, 60.356-000 Fortaleza, CE, Brazil. <sup>29</sup>Farming Systems Ecology, Wageningen University and Research, P.O. Box 430, 6700 AK Wageningen, Netherlands. <sup>30</sup>Chair of Nature Conservation and Landscape Ecology, University of Freiburg, Tennebacher Straße 4, 79106 Freiburg, Germany. <sup>31</sup>LTZER Zone Atelier Plaine and Val de Sevre, CEBC UMR 7372, CNRS and Université de La Rochelle, Beauvoir sur Niort 79360, France. <sup>32</sup>Arthropods Department, Natural Sciences Museum of Barcelona, 08003 Barcelona, Spain. <sup>33</sup>Instituto Nacional de Tecnología Agropecuaria, Estación Experimental Concordia, Estacion Yuqueri y vias del Ferrocarril s/n, 3200 Entre Rios, Argentina. <sup>34</sup>Faculty of Biology, University of Belgrade, Studentski trg 16, 11000 Belgrade, Serbia. <sup>35</sup>Instituto de Ecología Regional (IER), Universidad Nacional de Tucumán, CONICET, 4107 Yerba Buena, Tucumán, Argentina. <sup>36</sup>W.K. Kellogg Biological Station, Michigan State University, Hickory Corners, MI 49060, USA. <sup>37</sup>Federal Institute of Education, Science and Technology of Mato Grosso, Campus of Barra do Garças/MT, 78600-000, Brazil. <sup>38</sup>Center of Sustainable Development, University of Brasília (UnB)—Campus Universitário Darcy Ribeiro, Asa Norte, Brasília-DF 70910-900, Brazil. <sup>39</sup>Wageningen Environmental Research, Wageningen University and Research, P.O. Box 47, 6700 AA Wageningen, Netherlands. <sup>40</sup>Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime, Kent ME44TB, UK. <sup>41</sup>INRA, UR 1115, Plantes et Systèmes de culture Horticoles, 84000 Avignon, France. <sup>42</sup>Centre for Agri-Environmental Research, School of Agriculture, Policy and Development, Reading University, Reading RG6 6AR, UK. <sup>43</sup>Department of Entomology, University of Wisconsin, Madison, WI 53705, USA. <sup>44</sup>Instituto Nacional de Pesquisas da Amazônia (INPA), CEP 69.067-375 Manaus, Amazonas, Brazil. <sup>45</sup>Human-Environment Systems, Ecology, Evolution, and Behavior, Department of Biological Sciences, Boise State University, Boise, ID 83725, USA. <sup>46</sup>Department of Integrative Biology, University of Texas at Austin, 205 W 24th Street, 401 Biological Laboratories, Austin, TX 78712, USA. <sup>47</sup>Department of Biology and Environment, University of Haifa, Oranim, Tivon 36006, Israel. <sup>48</sup>Department of Entomology, Kansas State University, 125 Waters Hall, Manhattan, KS 66503, USA. <sup>49</sup>Department of Evolutionary and Environmental Biology, University of Haifa, 3498838 Haifa, Israel. <sup>50</sup>Institute for Plant Science and Microbiology, University of Hamburg, Hamburg, Germany. <sup>51</sup>Biodiversity International, Bangalore 560 065, India. <sup>52</sup>Ashoka Trust for Research in Ecology and the Environment (ATREE), Bangalore, India. <sup>53</sup>Bren School of Environmental Science and Management, University of California, Santa Barbara, Santa Barbara, CA 93106-5131, USA. <sup>54</sup>Department of Environmental Studies, Seattle University, 901 12th Avenue, Seattle, WA 9812, USA. <sup>55</sup>Department of Botany and Biodiversity Research, Division of Conservation Biology, Vegetation Ecology and Landscape Ecology, University of Vienna, Rennweg 14, 1030 Vienna, Austria. <sup>56</sup>Department of Entomology and Nematology, University of Florida, 1881 Natural Area Drive, Gainesville, FL 32601, USA. <sup>57</sup>Agrosavia, Centro de Investigación Obonuco, Km 5 via Obonuco, Pasto, Nariño, Colombia. <sup>58</sup>Agriculture, Livestock and Agroforestry Program, Tropical Agricultural Research and Higher Education Center (CATIE), Cartago, Turrialba 30501, Costa Rica. <sup>59</sup>National Audubon Society, Boulder, CO 80305, USA. <sup>60</sup>Institute for Resources, Environment and Sustainability, University of British Columbia, Vancouver, BC, Canada. <sup>61</sup>Cátedra de Avicultura, Cunicultura y Apicultura, Facultad de Agronomía, Universidad de Buenos Aires, CABA C1417DSE, Argentina. <sup>62</sup>School of Plant and Environmental Sciences, Virginia Tech, Blacksburg, VA 24061, USA. <sup>63</sup>Department of Ecology, UnB—Campus Universitário Darcy Ribeiro, Brasília-DF 70910-900, Brazil. <sup>64</sup>Department of Entomology and Nematology, University of California, Davis, Davis, CA 95616, USA. <sup>65</sup>INRA, UMR 1065 Santé et Agroécologie du Vignoble, ISVV, Université de Bordeaux, Bordeaux Sciences Agro, F-33883 Villenave d'Ornon Cedex, France. <sup>66</sup>INIBIOMA, Universidad Nacional del Comahue, CONICET, Quintral 1250, 8400 Bariloche, Río Negro, Argentina. <sup>67</sup>Senckenberg Biodiversity and Climate Research Centre (SBK-F), Senckenberganlage 25, 60325 Frankfurt am Main, Germany. <sup>68</sup>Centre for Biodiversity and Biosecurity, University of Auckland, Auckland, New Zealand. <sup>69</sup>Department of Environmental Science, Policy, and Management, University of California Berkeley, Berkeley, CA, USA. <sup>70</sup>South African National Biodiversity Institute, Kirstenbosch Research Centre, Private Bag X7, Claremont 7735, South Africa. <sup>71</sup>School of Agriculture and Food Science and Earth Institute, University College Dublin, Belfield, Dublin 4, Ireland. <sup>72</sup>School of Natural Sciences, Trinity College Dublin, Dublin 2, Ireland. <sup>73</sup>Institute for Sustainable Agro-ecosystem Services, School of Agriculture and Life Sciences, The University of Tokyo, 188-0002 Tokyo, Japan. <sup>74</sup>Forestry and Forest Products Research Institute, 1 Matsunosato, Tsukuba, Ibaraki 305-8687, Japan. <sup>75</sup>Instituto de Biologia, Universidade Federal da Bahia, 40170-210 Salvador, Brazil. <sup>76</sup>Bio-Protection Research Centre, Lincoln University, Lincoln, New Zealand. <sup>77</sup>Fukushima Branch, National Institute for Environmental Studies, 963-770 Fukushima, Japan. <sup>78</sup>Environment and Production Technology Division, International Food Policy Research Institute, Washington, DC 20005, USA. <sup>79</sup>Department of Health and Environmental Sciences, Xi'an Jiaotong-Liverpool University, 215123, Suzhou, China.

\*Corresponding author. Email: matteo.dainese@eurac.edu



**Fig. 1. Distribution of analyzed studies and effects of richness on ecosystem services provisioning.** (A) Map showing the size (number of crop fields sampled) and location of the 89 studies (further details of studies are given in table S1). (B) Global effect of pollinator richness on pollination ( $n = 821$  fields of 52 studies). (C) Global effect of natural enemy richness on pest control ( $n = 654$  fields of 37 studies). The thick line in each plot represents the median of the posterior distribution of the model. Light gray lines represent 1000 random draws from the posterior. The lines are included to depict uncertainty of the modeled relationship.

dominance) as the  $E_{var}$  index (27) that reflects the relative abundance of those taxa. We derived a standardized index of pollination services using measures of pollination success (fruit set or seed set) and plant reproduction (pollen limitation) and of pest control services using measures of natural enemy activity (predation or parasitism rates) and crop damage (pest activity) (see Materials and Methods). We  $z$ -transformed each measure separately to remove the effect of differences in measurement scale between indices and inverted values for those measures where low values indicate positive contribution to ecosystem service (i.e., pest activity). We also characterized the 1-km-radius landscape surrounding each field by measuring the percentage of cropland from high-resolution land-use maps. This landscape metric has been used as a relevant proxy for characterizing landscape simplification (28, 29) and is often correlated with other indicators of landscape complexity (30, 31).


Using a Bayesian multilevel modeling approach, we addressed three fundamental yet unresolved questions in the biodiversity-ecosystem function framework: (i) Do richness, abundance, and dominance sustain ecosystem services in real-world ecosystems? (ii) Does landscape simplification indirectly affect ecosystem services mediated by a loss

of local community diversity? (iii) How strong are the cascading effects of landscape simplification on final crop production?

## RESULTS AND DISCUSSION

We found clear evidence that richness of service-providing organisms positively influenced ecosystem service delivery. This was detected for both pollination (Fig. 1B and table S2) and pest control (Fig. 1C and table S2) and in almost all studies (figs. S1 and S2). As different methods were used in different studies to quantify richness and ecosystem services, we tested the sensitivity of our results to methodological differences. The bivariate relationships between richness and ecosystem services were robust to the taxonomic resolution to which each organism was identified, the sampling methods used to collect pollinators and natural enemies, and the pollination and pest control service measures used (see Materials and Methods).

The positive contribution of richness to service supply was also confirmed in a series of path analyses where we partitioned the relative importance of richness, total abundance, and evenness in driving biodiversity-ecosystem services relationships (Fig. 2). Although richness




**Fig. 2. Direct and indirect effects of richness, total abundance, and evenness on ecosystem services.** (A) Path model of pollinator richness as a predictor of pollination, mediated by pollinator abundance. (B) Path model of natural enemy richness as a predictor of pest control, mediated by natural enemy abundance. (C) Path model of pollinator richness as a predictor of pollination, mediated by pollinator evenness. (D) Path model of natural enemy richness as a predictor of pest control, mediated by natural enemy evenness. Pollination model,  $n = 821$  fields of 52 studies; pest control model,  $n = 654$  fields of 37 studies. Path coefficients are effect sizes estimated from the median of the posterior distribution of the model. Black and red arrows represent positive or negative effects, respectively. Arrow widths are proportional to highest density intervals (HDIs).

showed the larger contribution in both models, we also found a direct effect of total abundance (Fig. 2, A and B) and evenness (Fig. 2, C and D) on ecosystem service delivery. While both richness and total abundance showed a positive effect, evenness had a negative effect, suggesting that dominant species contribute more strongly to service supply. This integrative assessment of different aspects of community structure revealed a more multilayered relationship between biodiversity and ecosystem services than has been previously acknowledged. Our results complement previous findings for pollination (7, 21, 32) and pest control (33) and indicate that richness and total and relative abundance are not mutually exclusive but concurrently contribute to support these two key ecosystem services to agriculture. Hence, we find strong support for the role of species-rich communities in supporting pollination and pest control services.

Furthermore, we found that landscape simplification indirectly affected ecosystem services by reducing the richness of service-providing organisms. Roughly a third of the negative effects of landscape simplification on pollination were due to a loss in pollinator richness (Fig. 3A and table S5). This effect was even greater for pest control where natural enemy richness mediated about 50% of the total effect of landscape simplification (Fig. 3B and table S5). A similar pattern was also found considering abundance in addition to richness. In this case, landscape simplification indirectly affected ecosystem services by reducing both richness and abundance of service-providing organisms (fig. S4, A and B, and table S6). A consistent richness-mediated effect was also confirmed when we tested the direct and indirect effects of landscape simplification on ecosystem services via changes in both richness and evenness (fig. S4, C and D, and table S7). However, contrary to our ex-





**Fig. 3. Direct and indirect effects of landscape simplification on richness of service-providing organisms and associated ecosystem services.** (A) Path model of landscape simplification as a predictor of pollination, mediated by pollinator richness ( $n = 821$  fields of 52 studies). (B) Path model of landscape simplification as a predictor of pest control, mediated by natural enemy richness ( $n = 654$  fields of 37 studies). Path coefficients are effect sizes estimated from the median of the posterior distribution of the model. Black and red arrows represent positive and negative effects, respectively. Arrow widths are proportional to HDIs. Gray arrows represent nonsignificant effects (HDIs overlapped zero).


pectation, landscape simplification led to higher pollinator evenness. It is likely that landscape simplification has stronger effects on more specialized, rare species and shifts assemblages toward more evenly abundant, mobile, generalist species with a higher ability to use scattered resources in the landscape (34, 35). These findings, combined with the larger contribution of richness when tested in combination with evenness, suggest that not only dominant species but also relatively rare spe-

cies contributed to pollination services (32, 36). The effect of landscape simplification on ecosystem services was minimized when not considering the mediated effect of richness, especially for pest control. We did not find a direct landscape simplification effect on pest control [all highest density intervals (HDIs) overlapped zero; Fig. 3B and table S5]. Together, these results demonstrate strong negative indirect effects of landscape simplification on biodiversity-driven ecosystem services in agroecosystems and the importance of the richness and abundance of service-providing organisms in mediating these effects.

Last, for a subset of the data that had crop production information (674 fields of 42 studies), we found that the cascading effects of landscape simplification, mediated through richness and associated ecosystem services, led to lower crop production. This was detected for both pollination (Fig. 4A and table S8) and pest control (Fig. 4B and table S8). Specifically, landscape simplification reduced both pollinator and natural enemy richness, which had indirect consequences for pollination and pest control and, in turn, decreased crop production. For pest control, a positive link with crop production was detected in fields where the study area was not sprayed with insecticides during the course of the experiment (Fig. 4B) but not when considering all sites combined (with and without insecticide use; fig. S4). In sprayed areas, we did not find a pest control effect (all HDIs overlapped zero), probably because effects were masked by insecticide use (37, 38), indicating that insecticide use undermines the full potential of natural pest control. A positive link with crop production was detected, although measures used to estimate pest control (natural enemy and pest activity) were not direct components of crop production, as was the case of pollination measures (fruit or seed set). We also found an indirect influence of dominance (Fig. 4 and table S8) but not of abundance (fig. S4 and table S9) on crop production. However, the indirect dominance pathway was weaker than that via richness. An abundance effect may not have been detectable because of the lower sample size in these submodels with crop production. Although only available from a subset of the data, this result supports the hypothesis that the effects of landscape simplification can cascade up to reducing the final provisioning service of crop production.

Our findings suggest that some previously inconsistent responses of natural enemy abundance and activity to surrounding landscape composition (39) can be reconciled by considering richness in addition to abundance. Although richness and abundance are often correlated, their response to environmental variation can differ. This was evident in the path analysis showing a strong effect of landscape simplification on richness but only a marginal effect on abundance (fig. S3B). Moreover, effect sizes for natural enemy abundances in individual studies (fig. S5) showed similarly inconsistent responses to the previous synthesis (39). Likewise, results for natural enemy activity showed inconsistent responses to surrounding landscape composition in both syntheses [Fig. 3B and (39)].

Using an integrative model to assess key ecological theory, we demonstrate that the negative effects of landscape simplification on service supply and final crop production are primarily mediated by loss of species. We found strong evidence for positive biodiversity–ecosystem service relationships, highlighting that managing landscapes to enhance the richness of service-providing organisms (40, 41) is a promising pathway toward a more sustainable food production globally. In an era of rapid environmental changes, preserving biodiversity-driven services will consistently confer greater resilience to agroecosystems, such that we could expect improved crop production under a broader range of potential future conditions.



**Fig. 4. Direct and cascading effects of landscape simplification on final crop production via changes in richness, evenness, and ecosystem services.** (A) Path model representing direct and indirect effects of landscape simplification on final crop production through changes in pollinator richness, evenness, and pollination ( $n = 438$  fields of 27 studies). (B) Path model representing direct and indirect effects of landscape simplification on final crop production through changes in natural enemy richness, evenness, and pest control [only insecticide-free areas were considered in the model ( $n = 185$  fields of 14 studies)]. Path coefficients are effect sizes estimated from the median of the posterior distribution of the model. Black and red arrows represent positive and negative effects, respectively. Arrow widths are proportional to HDIs. Gray arrows represent nonsignificant effects (HDIs overlapped zero).

## MATERIALS AND METHODS

### Database compilation

We compiled data from crop studies where measures of richness and abundance of service-providing organisms (pollinators or natural enemies) and associated ecosystem services (pollination and biological

pest control) were available for the same sites. If available, we also included information on yield. Studies were identified by first searching the reference lists of recent meta-analyses (7, 21, 39, 42, 43) and then directly contacting researchers. For pest control, data were mostly provided from a recent pest control database (39). Of 191 researchers

initially contacted, 86 provided data that met our criteria. As similar studies were frequently performed in the same area, occasionally in the same year, and studies with multiple years usually used different sites each year, we did not nest year within study. Likewise, some studies collected data in different crops. Thus, we considered each year of multi-year studies (that is, 10 studies) and each crop of multicrop studies (that is, 5 studies) to be an independent dataset and used study-crop-year combinations as the highest hierarchical unit. Overall, we analyzed data from 89 studies and 1475 fields in 27 countries around the world (table S1). Twenty-nine crops were considered, including a wide array of annual and perennial fruit, seed, nut, stimulant, pulse, cereal, and oilseed crops. Studies represented the spectrum of management practices, including conventional, low-input conventional, integrated pest management, and organic farming. Few studies compared different management practices in similar landscapes, crop types, or sampled year. In 76% of fields, pest control experiments were performed in insecticide-free areas. In some fields, this information was not available (7%) or insecticides were applied (17%).

### Pollinator and pest natural enemy richness and abundance

Studies used a broad range of methods, which we categorized as active or passive (44), to sample pollinators or natural enemies. Active sampling methods included netting pollinators seen on crop flowers, hand-collecting individuals on plants, observational counting, sweep-netting, and vacuum sampling. Passive sampling methods were malaise traps, pan traps, pitfall traps, and sticky cards. Active sampling was performed in 85% of pollinator sampling fields and in 50% of natural enemy sampling fields.

Pollinators included representatives from the orders Hymenoptera, Diptera, Lepidoptera, and Coleoptera. Bees (Hymenoptera: Apoidea) were the most commonly observed pollinators and included *Apis* bees (Apidae: *Apis mellifera*, *Apis cerana*, *Apis dorsata*, and *Apis florea*), stingless bees (Apidae: Meliponini), bumblebees (Apidae: *Bombus* spp.), carpenter bees (Apidae: Xylocopini), small carpenter bees (Apidae: Ceratinini), sweat bees (Halictidae), long-horned bees (Apidae: Eucerini), plasterer bees (Colletidae), mining bees (Andrenidae), and mason bees (Megachilidae). Non-bee taxa included syrphid flies (Diptera: Syrphidae), other flies (Diptera: Calliphoridae, Tachinidae, and Muscidae), butterflies and moths (Lepidoptera), various beetle families (Coleoptera), and hymenopterans including ants (Formicidae) and the paraphyletic group of non-bee aculeate wasps. Natural enemies included ground beetles (Coleoptera), flies (Diptera), spiders (Aranea), hymenopterans including ants (Formicidae) and wasps, bugs (Hemiptera), thrips (Thysanoptera), net-winged insects (Neuroptera), bats, and birds.

We calculated pollinator and natural enemy richness as the number of unique taxa sampled per study, method, and field. A taxon was defined as a single biological type (that is, species, morphospecies, genus, and family) determined at the finest taxonomic resolution to which each organism was identified. In almost 70% of cases, taxonomic resolution was to species level (averaged proportion among all studies), but sometimes, it was based on morphospecies (15%), genus (8%), or family levels (7%). Taxon richness per field varied between 1 and 49 for pollinators and between 1 and 40 for natural enemies. Abundance reflected the sum of individuals sampled per study, method, and field. Pollinator richness was calculated either including or excluding honey bees (*A. mellifera*). *A. mellifera* was considered as the only species within the honey bee group for consistency across all datasets (43). Other *Apis* bees (that is, *A. cerana*, *A. dorsata*, and *A. florea*) were not pooled into the honey bee category as the large majority of observed individ-

uals were derived from feral populations. Feral and managed honey bees were analyzed as a single group because they cannot be distinguished during field observations. Feral honey bees were uncommon in most studies except for those in Africa and South America (*A. mellifera* is native in Africa, while it was introduced to the Americas). In studies with subsamples within a field (that is, plots within fields or multiple sampling rounds within fields), we calculated the total number of individuals and unique taxa across these subsamples.

We also calculated evenness using the  $E_{\text{var}}$  index (27). This index is based on the variance of log abundances (centered on the mean of log abundances) and then appropriately scaled to cover 0 to 1 (0, maximally uneven and 1, perfectly even).

### Pollination and pest control services

As different methods were used to quantify pollination or pest control services across studies, standardization was necessary to put all the indices on equivalent terms. Therefore, we transformed each index  $y$  in each field  $i$  in each study  $j$  using  $z$  scores. We preferred the use of  $z$  scores over other transformations (for example, division by the maximum) because  $z$  scores do not constrain the variability found in the raw data, as do other indices that are bounded between 0 and 1. We used the proportion of flowers that set fruit (that is, fruit set), the average number of seeds per fruits (that is, seed set), or the estimated measures of pollinator contribution to plant reproduction (that is, differences in fruit weight between plants with and without insect pollination, hereafter  $\Delta$  fruit weight) as measures of pollination services. We then converted these measures into the pollination index. The pest control index was calculated using measures of natural enemy activity or pest activity. Natural enemy activity was measured by sentinel pest experiments where pests were placed in crop fields and predation or parasitism rates were monitored or by field enclosure experiments where cages were used to exclude natural enemies to quantify differences in pest abundance or crop damage between plants with and without natural enemies. Pest activity was measured as the fraction or amount of each crop consumed, infested, or damaged. We inverted standardized values of pest activity by multiplying by  $-1$ , as low values indicate positive contributions to the ecosystem service.

### Crop production

Depending on the crop type, marketable crop yield is valued by farmers not only in terms of area-based yield but also in terms of fruit or seed weight [for example, in coffee, sunflower, or strawberry fields; (45, 46)] or seed production per plant [for example, in seed production fields; (36)]. Moreover, area-based yield and within-plant yield are often correlated (35, 36). Thus, we used both area-based yield and within-plant yield as measures of final crop production. Within-plant yield was measured by the total number (or mass) of seeds or fruits per plant or by fruit or seed weight. In addition, in this case, we standardized variables ( $z$  scores) to put all the indices on equivalent terms.

### Landscape simplification

Landscapes were characterized by calculating the percentage of cropland (annual and perennial) within a radius of 1 km around the center of each crop field. This landscape metric has been used as a relevant proxy for characterizing landscape simplification (28, 29) and is often correlated with other indicators of landscape complexity (30, 31). Moreover, we used this metric because cropland data are readily accessible from publicly available land cover data and are more accurate than other land use types such as forests and grasslands (47), especially when

detailed maps are not available. The 1-km spatial extent was chosen to reflect the typical flight and foraging distances of many insects including pollinators (48, 49) and natural enemies (50, 51). For studies where this information was not supplied by the authors, land uses were digitized using GlobeLand30 (52), a high-resolution map of Earth's land cover. The derived land cover maps were verified and, if necessary, corrected using a visual inspection of satellite images (Google Earth). We then calculated the percentage of cropland within the radius using Quantum GIS 2.18 (Open Source Geospatial Foundation Project; <http://qgis.osgeo.org>). The average percentage of cropland was 67.5% for pollination studies and 41.5% for natural enemy studies.

## Data analysis

### Data standardization

Before performing the analyses, we standardized the predictors (abundance, richness, and landscape simplification) using  $z$  scores within each study. This standardization was necessary to allow comparisons between studies with differences in methodology and landscape ranges (53). Moreover, this allows the separation of within-study effects from between-study effects. This separation is important because it prevents the risk of misinterpreting the results based on studies differing in methodology and landscape gradients, by erroneously extrapolating from between-study effects to within-study effects or vice versa (54, 55).

### Relationship between richness and ecosystem services

The relationship between richness of service-providing organisms and related ecosystem services (Fig. 1, B and C) was estimated from a Bayesian multilevel (partial pooling) model that allowed the intercept and the slope to vary among studies (also commonly referred to as random intercepts and slopes), following the equation

$$\begin{aligned} ES_i &\sim N(\alpha_{j[i]} + \beta_{j[i]} RIC_i, \sigma_j) \\ \alpha_j &\sim N(\mu_\alpha, \sigma_\alpha) \\ \beta_j &\sim N(\mu_\beta, \sigma_\beta) \end{aligned}$$

where  $ES_i$  is the ecosystem service index (pollination or pest control depending on the model),  $RIC_i$  is richness of service-providing organisms (pollinator or natural enemy richness depending on the model), and  $j_{[i]}$  represents observation  $i$  of study  $j$ . This partial-pooling model estimated both study-level responses [yielding an estimate for each study ( $\beta_j$ )] and the distribution from which the study-level estimates were drawn, yielding a higher-level estimate of the overall response across crop systems ( $\mu_\beta$ ). In addition, it accounted for variation in variance and sample size across observations (for example, studies). The intercepts  $\alpha_j$  and slopes  $\beta_j$  varied between studies according to a normal distribution with mean  $\mu$  and SD  $\sigma$ . Independent within-study errors also followed a normal distribution  $\varepsilon_i \sim N(0, \sigma)$ . We used weakly informative priors: normal (0, 10) for the population-level parameters ( $\alpha, \beta$ ) and half-Student  $t(3, 0, 5)$  for the group-level SD and residual SD.

### Direct and indirect effects of richness, abundance, and evenness on service provisioning

As natural communities vary not only in number of species but also in relative abundance of each species (evenness) and the total number of individuals (abundance), it is important to incorporate these attributes when assessing or modeling biodiversity effects (12, 56). In a Bayesian multivariate response model, a form of path analysis, we partitioned

the relative importance of richness and total and relative abundance in driving biodiversity-ecosystem relationships. We hypothesized that richness drives both abundance and evenness according to a revised version of the “more individuals hypothesis” (57) that proposes that more species can exploit more diverse resources and may therefore maintain more individuals than species-poor communities. Specifically, we tested (i) whether richness per se directly influences ecosystem services or is instead mediated by abundance (Eq. 1, 3; Fig. 2, A and B) and (ii) whether richness per se directly influences ecosystem services or is instead mediated by evenness (Eq. 2, 4; Fig. 2, C and D). Before analysis, we also checked for data collinearity among abundance, evenness, and richness by calculating the variance inflation factor (VIF). No signal of collinearity was detected in either model (VIFs were below 1.8). In a preliminary analysis, we also tested a possible interaction between richness and evenness. No significant interaction was found in both pollination and pest control models. To illustrate, we first show the univariate multilevel (partial pooling) models following these equations

$$RIC_i \sim N(\alpha_{j[i]} + \beta_{j[i]} ABU_i, \sigma_j) \quad (1)$$

$$RIC_i \sim N(\alpha_{j[i]} + \beta_{j[i]} EVE_i, \sigma_j) \quad (2)$$

$$ES_i \sim N(\alpha_{j[i]} + \beta_{j[i]} RIC_i + \beta_{j[i]} ABU_i, \sigma_j) \quad (3)$$

$$ES_i \sim N(\alpha_{j[i]} + \beta_{j[i]} RIC_i + \beta_{j[i]} EVE_i, \sigma_j) \quad (4)$$

where  $RIC_i$  is richness of service-providing organisms,  $ABU_i$  is abundance,  $EVE_i$  is evenness,  $ES_i$  is the ecosystem service index, and the index  $j_{[i]}$  represents observation  $i$  of study  $j$ . We then specified both multivariate multilevel models in a matrix-vector notion (53), as follows

$$\begin{aligned} Y_i &\sim N(X_i B_{r[i]}, \Sigma_j) \\ B_r &\sim N(M_B, \Sigma_B) \end{aligned}$$

where  $Y_i$  is the matrix of response variables with observations  $i$  as rows and variables  $r$  as columns,  $X_i$  is the matrix of all predictors for response  $r$ ,  $B_r$  are the regression parameters ( $\alpha$  and  $\beta$ ) for response  $r$ ,  $M_B$  represents the mean of the distribution of the regression parameters, and  $\Sigma_B$  is the covariance matrix representing the variation of the regression parameters in the population groups. We used weakly informative priors: normal (0, 10) for the population-level parameters ( $\alpha, \beta$ ) and half-Student  $t(3, 0, 5)$  for the group-level SD and residual SD. In building the model, we ensured that no residual correlation between  $ES_i$  and  $RIC_i$ , between  $ES_i$  and  $EVE_i$ , or between  $ES_i$  and  $ABU_i$  was estimated [see the “set\_rescor” function in the package *brms*; (58)].

### Direct and indirect effects of landscape simplification on ecosystem services

To estimate the direct and indirect effects of landscape simplification on richness and associated ecosystem services, we used two models. In a Bayesian multivariate response model with causal mediation effects (hereafter, mediation model), we tested whether landscape simplification directly influences ecosystem services or is mediated by richness. Mediation analysis is a statistical procedure to test whether the effect of an independent variable  $X$  on a dependent variable  $Y$  ( $X \rightarrow Y$ ) is at least partly explained via the inclusion of a third hypothetical



variable, the mediator variable  $M (X \rightarrow M \rightarrow Y)$  (59). The three causal paths  $a$ ,  $b$ , and  $c'$  correspond to  $X$ 's effect on  $M$ ,  $M$ 's effect on  $Y$ , and  $X$ 's effect on  $Y$  accounting for  $M$ , respectively. The three causal paths correspond to parameters from two regression models, one in which  $M$  is the outcome and  $X$  is the predictor and one in which  $Y$  is the outcome and  $X$  and  $M$  are the simultaneous predictors (fig. S6). From these parameters, we can compute the mediation effect (the product  $ab$ , also known as the indirect effect) and the total effect of  $X$  on  $Y$

$$c = c' + ab$$

Thus, the total causal effect of  $X$ , which is captured by the parameter  $c$ , can be decomposed precisely into two components, a direct effect  $c'$  and an indirect (mediation) effect  $ab$  (the product of paths  $a$  and  $b$ ). The model included the ecosystem service index as response, landscape simplification as predictor, and richness as mediator (Fig. 3). The separate regression models that made up the Bayesian multivariate multilevel model followed these equations

$$RIC_i \sim N(\alpha_{j[i]} + \beta_{j[i]}LAND_i, \sigma_j)$$

$$ES_i \sim N(\alpha_{j[i]} + \beta_{j[i]}RIC_i + \beta_{j[i]}LAND_i, \sigma_j)$$

We then compiled a multilevel path analysis testing the direct and indirect effects of landscape simplification on ecosystem services via changes in both richness and abundance (Eq. 5, 6, 8; fig. S4, A and B) or richness and evenness (Eq. 5, 7, 9; fig. S4, C and D). The separate regression models that made up the model followed these equations

$$RIC_i \sim N(\alpha_{j[i]} + \beta_{j[i]}LAND_i, \sigma_j) \tag{5}$$

$$ABU_i \sim N(\alpha_{j[i]} + \beta_{j[i]}LAND_i, \sigma_j) \tag{6}$$

$$EVE_i \sim N(\alpha_{j[i]} + \beta_{j[i]}LAND_i, \sigma_j) \tag{7}$$

$$ES_i \sim N(\alpha_{j[i]} + \beta_{j[i]}RIC_i + \beta_{j[i]}ABU_i + \beta_{j[i]}LAND_i, \sigma_j) \tag{8}$$

$$ES_i \sim N(\alpha_{j[i]} + \beta_{j[i]}RIC_i + \beta_{j[i]}EVE_i + \beta_{j[i]}LAND_i, \sigma_j) \tag{9}$$

where  $RIC_i$  is richness of service-providing organisms,  $LAND_i$  is landscape simplification measured as the percentage of arable land surrounding each study site,  $ABU_i$  is abundance,  $EVE_i$  is evenness,  $ES_i$  is the ecosystem service index, and the index  $j_{[i]}$  represents observation  $i$  of study  $j$ . We then specified multivariate multilevel models in a matrix-vector notion, as explained above. The mediation analysis was implemented using the R package sjstats [v. 0.15.0; (60)].

### Cascading effects of landscape simplification on final crop production

For 42 studies and 675 fields (pollination model,  $n = 438$  fields of 27 studies; pest control model,  $n = 236$  fields of 15 studies; table S1), the data allowed us to use a multilevel path analysis to examine cascading effects of landscape simplification on final crop produc-

tion via changes in richness, abundance, evenness, and ecosystem services. We expected that (i) landscape simplification would have a direct effect on richness, abundance, and dominance of service-providing organisms and (ii) richness and abundance dominance would relate positively to intermediate services, which, in turn, (iii) would increase final crop production (Fig. 4 and fig. S4). The indirect effects of richness and abundance (Eq. 10, 11, 13, 15; Fig. 4) or richness and evenness (Eq. 10, 12, 14, 15; fig. S4) were tested separately

$$RIC_i \sim N(\alpha_{j[i]} + \beta_{j[i]}LAND_i, \sigma_j) \tag{10}$$

$$ABU_i \sim N(\alpha_{j[i]} + \beta_{j[i]}LAND_i, \sigma_j) \tag{11}$$

$$EVE_i \sim N(\alpha_{j[i]} + \beta_{j[i]}LAND_i, \sigma_j) \tag{12}$$

$$ES_i \sim N(\alpha_{j[i]} + \beta_{j[i]}RIC_i + \beta_{j[i]}ABU_i, \sigma_j) \tag{13}$$

$$ES_i \sim N(\alpha_{j[i]} + \beta_{j[i]}RIC_i + \beta_{j[i]}EVE_i, \sigma_j) \tag{14}$$

$$PROD_i \sim N(\alpha_{j[i]} + \beta_{j[i]}ES_i, \sigma_j) \tag{15}$$

where  $RIC_i$  is richness of service-providing organisms,  $LAND_i$  is landscape simplification measured as the percentage of arable land surrounding each study site,  $ABU_i$  is abundance,  $EVE_i$  is evenness,  $ES_i$  is the ecosystem service index,  $PROD_i$  is crop production, and the index  $j_{[i]}$  represents observation  $i$  of study  $j$ . We specified a multivariate multilevel model in a matrix-vector notion, as explained above.

### Parameter estimation

All analyses were conducted in the programming language Stan through R (v. 3.4.3) using the package brms [v. 2.2.0; (58)]. Stan implements Hamiltonian Monte Carlo and its extension, the No-U-Turn Sampler. These algorithms converge much more quickly than other Markov chain Monte Carlo algorithms especially for high-dimensional models (61). Each model was run with four independent Markov chains of 5000 iterations, discarding the first 2500 iterations per chain as warm-up and resulting in 10,000 posterior samples overall. Convergence of the four chains and sufficient sampling of posterior distributions were confirmed by (i) the visual inspection of parameter traces, (ii) ensuring a scale reduction factor ( $\bar{R}$ ) below 1.01, and (iii) an effective size ( $n_{\text{eff}}$ ) of at least 10% of the number of iterations. For each model, posterior samples were summarized on the basis of the Bayesian point estimate (median), SE (median absolute deviation), and posterior uncertainty intervals by HDIs, a type of credible interval that contains the required mass such that all points within the interval have a higher probability density than points outside the interval (62). The advantage of the Bayesian approach is the possibility to estimate not only expected values for each parameter but also the uncertainty associated with these estimates (63). Thus, we calculated 80, 90, and 95% HDIs for parameter estimates.

### Sensitivity analyses

Given that different methods were used in different studies to quantify richness, ecosystem services, and final crop production, we measured the sensitivity of our results to methodological differences.

(1) We verified whether treating each annual dataset from multiyear studies separately could incorrectly account for the dependence of the

data. We refitted the model testing the relationship between richness and ecosystem services including year nested within study (that is, study defined as study-crop combination). Then, we compared models (year-independent model versus year-nested model) using leave-one-out (LOO) cross-validation, a fully Bayesian model selection procedure for estimating pointwise out-of-sample predictive accuracy (64). We calculated the expected log pointwise predictive density ( $\widehat{\text{elpd}}_{100}$ ), using the log-likelihood evaluated at the posterior simulations of the parameter values. Model comparison was implemented using R package loo [v. 2.0.0; (65)]. We found that the year-nested model had a lower average predictive accuracy than the year-independent model for both pollination ( $\Delta\widehat{\text{elpd}}_{100} = -1.79$ ) and pest control ( $\Delta\widehat{\text{elpd}}_{100} = -1.09$ ) and therefore retained the year-independent model in our analysis.

(2) We verified whether taxonomic resolution influenced the interpretation of results. We recalculated richness considering only organisms classified at the fine taxonomy level (species or morphospecies levels) and refitted the model testing the effect of richness on ecosystem services. We found no evidence that taxonomic resolution influenced our results. With a fine taxonomic resolution, the effects of richness on ecosystem services ( $\beta_{\text{pollinators}} = 0.1535$ , 90% HDIs = 0.0967 to 0.2141;  $\beta_{\text{enemies}} = 0.2264$ , 90% HDIs = 0.1475 to 0.3065; table S2) were nearly identical to the estimates presented in the main text ( $\beta_{\text{pollinators}} = 0.1532$ , 90% HDIs = 0.0892 to 0.2058;  $\beta_{\text{enemies}} = 0.2093$ , 90% HDIs = 0.1451 to 0.2779; table S2).

(3) We verified whether the sampling methods used to collect pollinators (active versus passive sampling techniques) influenced the relationship between pollinator richness and pollination using Bayesian hypothesis testing (58). Passive methods do not directly capture flower visitors and may introduce some bias (for example, they may underestimate flower visitors). However, our estimate was not influenced by sampling method (the one-sided 90% credibility interval overlapped zero; table S11). In accordance with this finding, the evidence ratio showed that the hypothesis tested (that is, estimates of studies with active sampling > estimates of studies with passive sampling) was only 0.78 times more likely than the alternative hypothesis.

(4) We verified whether methodological differences in measuring pollination and pest control services influenced the relationship between richness and ecosystem services. Using Bayesian hypothesis testing, we tested whether the estimates differed among methods. The two-sided 95% credibility interval overlapped zero in all comparisons (estimates did not differ significantly; table S12), indicating that our estimate was not influenced by methodological differences in measuring ecosystem services. Furthermore, we tested effects including only inverted pest activity as a reflection of pest control. We found positive effects of natural enemy richness on inverted pest activity ( $\beta = 0.1307$ , 90% HDIs = 0.0102 to 0.2456), indicating that results were robust to the type of pest control measure considered.

(5) As honey bees are the most important and abundant flower visitors in some locations, we verified the potential influence of honey bees on our results by refitting the path models testing direct and indirect effects of richness, abundance, and evenness on pollination services with honey bees. A positive direct contribution of richness to pollination was confirmed even after including honey bees (fig. S7). In these models, both abundance and evenness showed a larger effect compared to models without honey bees (Fig. 2).

(6) Insecticide application during the course of the experiment could mask the effect of pest control on crop production (37, 38). We verified the potential influence of insecticide application on our results by refit-

ting the model considering only fields where the study area was not sprayed with insecticide during the course of the experiment ( $n = 85$  fields of 14 studies). We found a pest control effect that was masked when considering all sites combined (with and without insecticide; fig. S4). We therefore show the insecticide-free model in the main text (Fig. 4).

(7) We verified the consistency of our results considering only studies that measured area-based yield (submodel). Only the cascading effects of landscape simplification on final crop production via changes in richness were tested in a simplified model. We found no evident differences between the submodel (fig. S8) and the full model presented in the main text (Fig. 4).

## SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <http://advances.sciencemag.org/cgi/content/full/5/10/eaax0121/DC1>

Supplementary Text

- Fig. S1. Forest plot of the effect of pollinator richness on pollination for individual studies.  
 Fig. S2. Forest plot of the effect of natural enemy richness on pest control for individual studies.  
 Fig. S3. Direct and indirect landscape simplification effects on ecosystem services via changes in richness and abundance or richness and evenness.  
 Fig. S4. Direct and cascading landscape simplification effects on final crop production via changes in natural enemy richness, abundance evenness, and pest control (all sites together, with and without insecticide application).  
 Fig. S5. Forest plot of the effect of landscape simplification on natural enemy abundance for individual studies.  
 Fig. S6. Mediation model.  
 Fig. S7. Direct and indirect effects of pollinator richness, abundance, and evenness (with honey bees) on pollination.  
 Fig. S8. Direct and cascading landscape simplification effects on area-based yield via changes in richness, and ecosystem services.  
 Table S1. List of 89 studies considered in our analyses.  
 Table S2. Model output for richness-ecosystem service relationships.  
 Table S3. Model output for path models testing direct and indirect effects (mediated by changes in abundance) of richness on ecosystem services.  
 Table S4. Model output for path models testing direct and indirect effects (mediated by changes in evenness) of richness on ecosystem services.  
 Table S5. Model output for path models testing direct and indirect effects (mediated by changes in richness) of landscape simplification on ecosystem services.  
 Table S6. Model output for path models testing the direct and cascading landscape simplification effects on ecosystem services via changes in richness and abundance.  
 Table S7. Model output for path models testing the direct and cascading landscape simplification effects on ecosystem services via changes in richness and evenness.  
 Table S8. Model output for path models testing the direct and cascading landscape simplification effects on final crop production via changes in richness, evenness, and ecosystem services.  
 Table S9. Model output for path models testing the direct and cascading landscape simplification effects on final crop production via changes in richness, abundance, and ecosystem services.  
 Table S10. Model output for path models testing direct and indirect effects of pollinator richness, abundance, and evenness (with honey bees) on pollination.  
 Table S11. Results of pairwise comparison of richness-ecosystem service relationships according to the methods used to sample pollinators and natural enemies.  
 Table S12. Results of pairwise comparison of richness-ecosystem service relationships according to the methods used to quantify pollination and pest control services.  
 Database S1. Data on pollinator and natural enemy diversity and associated ecosystem services compiled from 89 studies and 1475 locations around the world.  
 References (66–111)

[View/request a protocol for this paper from Bio-protocol.](#)

## REFERENCES AND NOTES

- S. Díaz, U. Pascual, M. Stenseke, B. Martín-López, R. T. Watson, Z. Molnár, R. Hill, K. M. A. Chan, I. A. Baste, K. A. Brauman, S. Polasky, A. Church, M. Lonsdale, A. Larigauderie, P. W. Leadley, A. P. E. van Oudenhoven, F. van der Plaats, M. Schröter, S. Lavorel, Y. Aumeeruddy-Thomas, E. Bukvareva, K. Davies, S. Demissew, G. Erpul, P. Failler, C. A. Guerra, C. L. Hewitt, H. Keune, S. Lindley, Y. Shirayama, Assessing nature's contributions to people. *Science* **359**, 270–272 (2018).

2. S. Díaz, J. Settele, E. Brondizio, Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (United Nations Paris, Fr., 2019), 1–39.
3. S. Naeem, J. E. Duffy, E. Zavaleta, The functions of biological diversity in an age of extinction. *Science* **336**, 1401–1406 (2012).
4. D. Tilman, F. Isbell, J. M. Cowles, Biodiversity and ecosystem functioning. *Annu. Rev. Ecol. Syst.* **45**, 471–493 (2014).
5. B. J. Cardinale, J. E. Duffy, A. Gonzalez, D. U. Hooper, C. Perrings, P. Venail, A. Narwani, G. M. Mace, D. Tilman, D. A. Wardle, A. P. Kinzig, G. C. Daily, M. Loreau, J. B. Grace, A. Larigauderie, D. S. Srivastava, S. Naeem, Biodiversity loss and its impact on humanity. *Nature* **486**, 59–67 (2012).
6. R. Winfree, J. W. Fox, N. M. Williams, J. R. Reilly, D. P. Cariveau, Abundance of common species, not species richness, drives delivery of a real-world ecosystem service. *Ecol. Lett.* **18**, 626–635 (2015).
7. D. Kleijn, R. Winfree, I. Bartomeus, L. G. Carvalheiro, M. Henry, R. Isaacs, A.-M. Klein, C. Kremen, L. K. M'Gonigle, R. Rader, T. H. Ricketts, N. M. Williams, N. Lee Adamson, J. S. Ascher, A. Báldi, P. Batáry, F. Benjamin, J. C. Biesmeijer, E. J. Blitzer, R. Bommarco, M. R. Brand, V. Bretagnolle, L. Button, D. P. Cariveau, R. Chifflet, J. F. Colville, B. N. Danforth, E. Elle, M. P. D. Garratt, F. Herzog, A. Holzschuh, B. G. Howlett, F. Jauker, S. Jha, E. Knop, K. M. Krewenka, V. le Féon, Y. Mandelik, E. A. May, M. G. Park, G. Pisanty, M. Reemer, V. Riedinger, O. Rollin, M. Rundlöf, H. S. Sardiñas, J. Scheper, A. R. Sciligo, H. G. Smith, I. Steffan-Dewenter, R. Thorp, T. Tscharntke, J. Verhulst, B. F. Viana, B. E. Vaissière, R. Veldtman, K. L. Ward, C. Westphal, S. G. Potts, Delivery of crop pollination services is an insufficient argument for wild pollinator conservation. *Nat. Commun.* **6**, 7414 (2015).
8. M. Loreau, A. Hector, Partitioning selection and complementarity in biodiversity experiments. *Nature* **412**, 72–76 (2001).
9. D. U. Hooper, F. S. Chapin III, J. J. Ewel, A. Hector, P. Inchausti, S. Lavorel, J. H. Lawton, D. M. Lodge, M. Loreau, S. Naeem, B. Schmid, H. Setälä, A. J. Symstad, J. Vandermeer, D. A. Wardle, Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecol. Monogr.* **75**, 3–35 (2005).
10. M. Schleuning, J. Fründ, D. García, Predicting ecosystem functions from biodiversity and mutualistic networks: An extension of trait-based concepts to plant-animal interactions. *Ecography* **38**, 380–392 (2015).
11. T. H. Larsen, N. M. Williams, C. Kremen, Extinction order and altered community structure rapidly disrupt ecosystem functioning. *Ecol. Lett.* **8**, 538–547 (2005).
12. J. Reiss, J. R. Bridle, J. M. Montoya, G. Woodward, Emerging horizons in biodiversity and ecosystem functioning research. *Trends Ecol. Evol.* **24**, 505–514 (2009).
13. K. H. Bannar-Martin, C. T. Kremer, S. K. M. Ernest, M. A. Leibold, H. Auge, J. Chase, S. A. J. Declerck, N. Eisenhauer, S. Harpole, H. Hillebrand, F. Isbell, T. Koffel, S. Larsen, A. Narwani, J. S. Petermann, C. Roscher, J. S. Cabral, S. R. Supp, Integrating community assembly and biodiversity to better understand ecosystem function: The community assembly and the functioning of ecosystems (CAFE) approach. *Ecol. Lett.* **21**, 167–180 (2018).
14. P. Balvanera, I. Siddique, L. Dee, A. Paquette, F. Isbell, A. Gonzalez, J. Byrnes, M. I. O'Connor, B. A. Hungate, J. N. Griffin, Linking biodiversity and ecosystem services: Current uncertainties and the necessary next steps. *Bioscience* **64**, 49–57 (2014).
15. F. Isbell, A. Gonzalez, M. Loreau, J. Cowles, S. Díaz, A. Hector, G. M. Mace, D. A. Wardle, M. I. O'Connor, J. E. Duffy, L. A. Turnbull, P. L. Thompson, A. Larigauderie, Linking the influence and dependence of people on biodiversity across scales. *Nature* **546**, 65–72 (2017).
16. R. Chaplin-Kramer, M. E. O'Rourke, E. J. Blitzer, C. Kremen, A meta-analysis of crop pest and natural enemy response to landscape complexity. *Ecol. Lett.* **14**, 922–932 (2011).
17. C. M. Kennedy, E. Lonsdorf, M. C. Neel, N. M. Williams, T. H. Ricketts, R. Winfree, R. Bommarco, C. Brittain, A. L. Burley, D. Cariveau, L. G. Carvalheiro, N. P. Chacoff, S. A. Cunningham, B. N. Danforth, J.-H. Dudenhöffer, E. Elle, H. R. Gaines, L. A. Garibaldi, C. Gratton, A. Holzschuh, R. Isaacs, S. K. Javorek, S. Jha, A. M. Klein, K. Krewenka, Y. Mandelik, M. M. Mayfield, L. Morandin, L. A. Neame, M. Otieno, M. Park, S. G. Potts, M. Rundlöf, A. Saez, I. Steffan-Dewenter, H. Taki, B. F. Viana, C. Westphal, J. K. Wilson, S. S. Greenleaf, C. Kremen, A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecol. Lett.* **16**, 584–599 (2013).
18. J. A. Foley, R. DeFries, G. P. Asner, C. Barford, G. Bonan, S. R. Carpenter, F. S. Chapin, M. T. Coe, G. C. Daily, H. K. Gibbs, J. H. Helkowski, T. Holloway, E. A. Howard, C. J. Kucharik, C. Monfreda, J. A. Patz, I. C. Prentice, N. Ramankutty, P. K. Snyder, Global consequences of land use. *Science* **309**, 570–574 (2005).
19. T. H. Ricketts, J. Regetz, I. Steffan-Dewenter, S. A. Cunningham, C. Kremen, A. Bogdanski, B. Gemmill-Herren, S. S. Greenleaf, A. M. Klein, M. M. Mayfield, L. A. Morandin, A. Ochieng', S. G. Potts, B. F. Viana, Landscape effects on crop pollination services: Are there general patterns? *Ecol. Lett.* **11**, 499–515 (2008).
20. F. J. J. A. Bianchi, C. J. H. Booij, T. Tscharntke, Sustainable pest regulation in agricultural landscapes: A review on landscape composition, biodiversity and natural pest control. *Proc. Biol. Sci.* **273**, 1715–1727 (2006).
21. L. A. Garibaldi, I. Steffan-Dewenter, R. Winfree, M. A. Aizen, R. Bommarco, S. A. Cunningham, C. Kremen, L. G. Carvalheiro, L. D. Harder, O. Afik, I. Bartomeus, F. Benjamin, V. Boreux, D. Cariveau, N. P. Chacoff, J. H. Dudenhöffer, B. M. Freitas, J. Ghazoul, S. Greenleaf, J. Hipolito, A. Holzschuh, B. Howlett, R. Isaacs, S. K. Javorek, C. M. Kennedy, K. M. Krewenka, S. Krishnan, Y. Mandelik, M. M. Mayfield, I. Motzke, T. Munyuli, B. A. Nault, M. Otieno, J. Petersen, G. Pisanty, S. G. Potts, R. Rader, T. H. Ricketts, M. Rundlöf, C. L. Seymour, C. Schuepp, H. Szentgyorgyi, H. Taki, T. Tscharntke, C. H. Vergara, B. F. Viana, T. C. Wagner, C. Westphal, N. Williams, A. M. Klein, Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science* **339**, 1608–1611 (2013).
22. D. K. Ray, N. Ramankutty, N. D. Mueller, P. C. West, J. A. Foley, Recent patterns of crop yield growth and stagnation. *Nat. Commun.* **3**, 1293 (2012).
23. J. A. Foley, N. Ramankutty, K. A. Brauman, E. S. Cassidy, J. S. Gerber, M. Johnston, N. D. Mueller, C. O'Connell, D. K. Ray, P. C. West, C. Balzer, E. M. Bennett, S. R. Carpenter, J. Hill, C. Monfreda, S. Polasky, J. Rockström, J. Sheehan, S. Siebert, D. Tilman, D. P. M. Zaks, Solutions for a cultivated planet. *Nature* **478**, 337–342 (2011).
24. R. Bommarco, D. Kleijn, S. G. Potts, Ecological intensification: Harnessing ecosystem services for food security. *Trends Ecol. Evol.* **28**, 230–238 (2013).
25. J. Pretty, Intensification for redesigned and sustainable agricultural systems. *Science* **362**, eaav0294 (2018).
26. Millennium Ecosystem Assessment, *Ecosystems and Human Well-being: Biodiversity Synthesis* (World Resources Institute, 2005).
27. B. Smith, J. B. Wilson, A consumer's guide to evenness indices. *Oikos* **76**, 70–82 (1996).
28. T. D. Meehan, B. P. Werling, D. A. Landis, C. Gratton, Agricultural landscape simplification and insecticide use in the Midwestern United States. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 11500–11505 (2011).
29. M. Dainese, N. J. B. Isaac, G. D. Powney, R. Bommarco, E. Öckinger, M. Kuussaari, J. Pöyry, T. G. Benton, D. Gabriel, J. A. Hodgson, W. E. Kunin, R. Lindborg, S. M. Sait, L. Marini, Landscape simplification weakens the association between terrestrial producer and consumer diversity in Europe. *Glob. Chang. Biol.* **23**, 3040–3051 (2017).
30. T. Tscharntke, A. M. Klein, A. Ruess, I. Steffan-Dewenter, C. Thies, Landscape perspectives on agricultural intensification and biodiversity—Ecosystem service management. *Ecol. Lett.* **8**, 857–874 (2005).
31. D. A. Landis, Designing agricultural landscapes for biodiversity-based ecosystem services. *Basic Appl. Ecol.* **18**, 1–12 (2017).
32. R. Winfree, J. R. Reilly, I. Bartomeus, D. P. Cariveau, N. M. Williams, J. Gibbs, Species turnover promotes the importance of bee diversity for crop pollination at regional scales. *Science* **359**, 791–793 (2018).
33. D. K. Letourneau, J. A. Jedlicka, S. G. Bothwell, C. R. Moreno, Effects of natural enemy biodiversity on the suppression of arthropod herbivores in terrestrial ecosystems. *Annu. Rev. Ecol. Syst.* **40**, 573–592 (2009).
34. L. Marini, E. Öckinger, K.-O. Bergman, B. Jauker, J. Krauss, M. Kuussaari, J. Pöyry, H. G. Smith, I. Steffan-Dewenter, R. Bommarco, Contrasting effects of habitat area and connectivity on evenness of pollinator communities. *Ecography* **37**, 544–551 (2014).
35. S. Gámez-Virués, D. J. Perović, M. M. Gossner, C. Börschig, N. Blüthgen, H. de Jong, N. K. Simons, A.-M. Klein, J. Krauss, G. Maier, C. Scherber, J. Steckel, C. Rothenwöhler, I. Steffan-Dewenter, C. N. Weiner, W. Weisser, M. Werner, T. Tscharntke, C. Westphal, Landscape simplification filters species traits and drives biotic homogenization. *Nat. Commun.* **6**, 8568 (2015).
36. T. P. M. Fijen, J. A. Scheper, T. M. Boom, N. Janssen, I. Raemakers, D. Kleijn, Insect pollination is at least as important for marketable crop yield as plant quality in a seed crop. *Ecol. Lett.* **21**, 1704–1713 (2018).
37. H. Liere, T. N. Kim, B. P. Werling, T. D. Meehan, D. A. Landis, C. Gratton, Trophic cascades in agricultural landscapes: Indirect effects of landscape composition on crop yield. *Ecol. Appl.* **25**, 652–661 (2015).
38. V. Gagic, D. Kleijn, A. Báldi, G. Boros, H. B. Jørgensen, Z. Elek, M. P. D. Garratt, G. A. de Groot, K. Hedlund, A. Kovács-Hostyánszki, L. Marini, E. Martin, I. Peveré, S. G. Potts, S. Redlich, D. Senapathi, I. Steffan-Dewenter, S. Switek, H. G. Smith, V. Takács, P. Tryjanowski, W. H. van der Putten, S. van Gils, R. Bommarco, Combined effects of agrochemicals and ecosystem services on crop yield across Europe. *Ecol. Lett.* **20**, 1427–1436 (2017).
39. D. S. Karp, R. Chaplin-Kramer, T. D. Meehan, E. A. Martin, F. DeClerck, H. Grab, C. Gratton, L. Hunt, A. E. Larsen, A. Martínez-Salinas, M. E. O'Rourke, A. Rusch, K. Poveda, M. Jonsson, J. A. Rosenheim, N. A. Schellhorn, T. Tscharntke, S. D. Wratten, W. Zhang, A. L. Iverson, L. S. Adler, M. Albrecht, A. Alignier, G. M. Angelella, M. Z. Anjum, J. Avelino, P. Batáry, J. M. Baveco, F. J. J. A. Bianchi, K. Birkhofer, E. W. Bohnenblust, R. Bommarco, M. J. Brewer, B. Caballero-López, Y. Carrière, L. G. Carvalheiro, L. Cayuela, M. Centrella, A. Četković, D. C. Henri, A. Chabert, A. C. Costamagna, A. De la Mora, J. de Kraker, N. Desneux, E. Diehl, T. Diekötter, C. F. Dormann, J. O. Eckberg, M. H. Entling, D. Fiedler, P. Franck, F. J. F. van Veen,



- T. Frank, V. Gagic, M. P. D. Garratt, A. Getachew, D. J. Gonthier, P. B. Goodell, I. Graziosi, R. L. Groves, G. M. Gurr, Z. Hajjan-Forooshani, G. E. Heimpel, J. D. Herrmann, A. S. Hueseth, D. J. Inčlán, A. J. Ingrao, P. Iv, K. Jacot, G. A. Johnson, L. Jones, M. Kaiser, J. M. Kaser, T. Keasar, T. N. Kim, M. Kishinevsky, D. A. Landis, B. Lavandero, C. Lavigne, A. Le Ralec, D. Lemessa, D. K. Letourneau, H. Liere, Y. Lu, Y. Lubin, T. Luttermoser, B. Maas, K. Mace, F. Madeira, V. Mader, A. M. Cortesero, L. Marini, E. Martinez, H. M. Martinson, P. Menozzi, M. G. E. Mitchell, T. Miyashita, G. A. R. Molina, M. A. Molina-Montenegro, M. E. O'Neal, I. Opatovskiy, S. Ortiz-Martinez, M. Nash, Ö. Östman, A. Ouin, D. Pak, D. Paredes, S. Parsa, H. Parry, R. Perez-Alvarez, D. J. Perović, J. A. Peterson, S. Petit, S. M. Philpott, M. Plantegenest, M. Plečaš, T. Pluess, X. Pons, S. G. Potts, R. F. Pywell, D. W. Ragsdale, T. A. Rand, L. Raymond, B. Ricci, C. Sargent, J. P. Sarthou, J. Saulais, J. Schächermann, N. P. Schmidt, G. Schneider, C. Schüepp, F. S. Sivakoff, H. G. Smith, K. S. Whitney, S. Stutz, Z. Szendrei, M. B. Takada, H. Taki, G. Tamburini, L. J. Thomson, Y. Tricault, N. Tsafack, M. Tschumi, M. Valantin-Morison, M. Van Trinh, W. van der Werf, K. T. Vierling, K. P. Werling, B. J. Wickens, V. J. Wickens, B. A. Woodcock, K. Wychuys, H. Xiao, M. Yasuda, A. Yoshioka, Y. Zou, Crop pests and predators exhibit inconsistent responses to surrounding landscape composition. *Proc. Natl. Acad. Sci. U. S. A.* **15**, E7863–E7870 (2018).
40. C. Kremen, A. M. Merenlender, Landscapes that work for biodiversity and people. *Science* **362**, eaau6020 (2018).
41. E. A. Martin, M. Dainese, Y. Clough, A. Baldi, R. Bommarco, V. Gagic, M. P. D. Garratt, A. Holzschuh, D. Kleijn, A. Kovács-Hostyánszki, L. Marini, S. G. Potts, H. G. Smith, D. Al Hassan, M. Albrecht, G. K. S. Andersson, J. D. Asís, S. Aviron, M. V. Balzan, L. Baños-Picón, I. Bartomeus, P. Batáry, F. Burel, B. Caballero-López, E. D. Concepción, V. Coudrain, J. Dänhardt, M. Diaz, T. Diekötter, C. F. Dormann, R. Duffot, M. H. Entling, N. Farwig, C. Fischer, T. Frank, L. A. Garibaldi, J. Hermann, F. Herzog, D. Inčlán, K. Jacot, F. Jauker, P. Jeanneret, M. Kaiser, J. Krauss, V. Le Féon, J. Marshall, A.-C. Moonen, G. Moreno, V. Riedinger, M. Rundlöf, A. Rusch, J. Scheper, G. Schneider, C. Schüepp, S. Stutz, L. Sutter, G. Tamburini, C. Thies, J. Tormos, T. Tschamtkke, M. Tschumi, D. Uzman, C. Wagner, M. Zubair-Anjum, I. Steffan-Dewenter, The interplay of landscape composition and configuration: New pathways to manage functional biodiversity and agroecosystem services across Europe. *Ecol. Lett.* **22**, 1083–1094 (2019).
42. L. A. Garibaldi, L. G. Carvalheiro, B. E. Vaissiere, B. Gemmill-Herren, J. Hipolito, B. M. Freitas, H. T. Ngo, N. Azzu, A. Saez, J. Astrom, J. An, B. Blochtein, D. Buchori, F. J. C. Garcia, F. Oliveira da Silva, K. Devkota, M. de Fátima Ribeiro, L. Freitas, M. C. Gaglianone, M. Goss, M. Irshad, M. Kasina, A. J. S. P. Filho, L. H. P. Kiill, P. Kwapong, G. N. Parra, C. Pires, V. Pires, R. S. Rawal, A. Rizali, A. M. Saraiva, R. Veldtman, B. F. Viana, S. Witter, H. Zhang, Mutually beneficial pollinator diversity and crop yield outcomes in small and large farms. *Science* **351**, 388–391 (2016).
43. R. Rader, I. Bartomeus, L. A. Garibaldi, M. P. D. Garratt, B. G. Howlett, R. Winfree, S. A. Cunningham, M. M. Mayfield, A. D. Arthur, G. K. S. Andersson, R. Bommarco, C. Brittain, L. G. Carvalheiro, N. P. Chacoff, M. H. Entling, B. Fouly, B. M. Freitas, B. Gemmill-Herren, J. Ghazoul, S. R. Griffin, C. L. Gross, L. Herbertsson, F. Herzog, J. Hipólito, S. Jaggard, F. Jauker, A. M. Klein, D. Kleijn, S. Krishnan, C. Q. Lemos, S. A. M. Lindström, Y. Mandelik, V. M. Monteiro, W. Nelson, L. Nilsson, D. E. Pattemore, N. de O. Pereira, G. Pisanty, S. G. Potts, M. Reemer, M. Rundlöf, C. S. Sheffield, J. Scheper, C. Schüepp, H. G. Smith, D. A. Stanley, J. C. Stout, H. Szentgyörgyi, H. Taki, C. H. Vergara, B. F. Viana, M. Woyciechowski, Non-bee insects are important contributors to global crop pollination. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 146–151 (2016).
44. E. M. Lichtenberg, C. M. Kennedy, C. Kremen, P. Batáry, F. Berendse, R. Bommarco, N. A. Bosque-Pérez, L. G. Carvalheiro, W. E. Snyder, N. M. Williams, R. Winfree, B. K. Klatt, S. Åström, F. Benjamin, C. Brittain, R. Chaplin-Kramer, Y. Clough, B. Danforth, T. Diekötter, S. D. Eigenbrode, J. Ekroos, E. Elle, B. M. Freitas, Y. Fukuda, H. R. Gaines-Day, H. Grab, C. Gratton, A. Holzschuh, R. Isaacs, M. Isaia, S. Jha, D. Jonason, V. P. Jones, A. M. Klein, J. Krauss, D. K. Letourneau, S. Macfadyen, R. E. Mallinger, E. A. Martin, E. Martinez, J. Memmott, L. Morandin, L. Neame, M. Otieno, M. G. Park, L. Pfiffner, M. J. O. Pocock, C. Ponce, S. G. Potts, K. Poveda, M. Ramos, J. A. Rosenheim, M. Rundlöf, H. Sardiñas, M. E. Saunders, N. L. Schon, A. R. Scilligo, C. S. Sidhu, I. Steffan-Dewenter, T. Tschamtkke, M. Vesely, W. W. Weisser, J. K. Wilson, D. W. Crowder, A global synthesis of the effects of diversified farming systems on arthropod diversity within fields and across agricultural landscapes. *Glob. Chang. Biol.* **23**, 4946–4957 (2017).
45. L. G. Carvalheiro, R. Veldtman, A. G. Shenkute, G. B. Tesfay, C. W. W. Pirk, J. S. Donaldson, S. W. Nicolson, Natural and within-farmland biodiversity enhances crop productivity. *Ecol. Lett.* **14**, 251–259 (2011).
46. V. Boreux, C. G. Kusalappa, P. Vaast, J. Ghazoul, Interactive effects among ecosystem services and management practices on crop production: Pollination in coffee agroforestry systems. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 8387–8392 (2013).
47. R. Tropek, O. Sedláček, J. Beck, P. Keil, Z. Musilova, I. Šimova, D. Storch, Comment on “High-resolution global maps of 21st-century forest cover change”. *Science* **344**, 981 (2014).
48. M. Rundlöf, J. Bengtsson, H. G. Smith, Local and landscape effects of organic farming on butterfly species richness and abundance. *J. Appl. Ecol.* **45**, 813–820 (2008).
49. A. Holzschuh, M. Dainese, J. P. González-Varo, S. Mudri-Stojnić, V. Riedinger, M. Rundlöf, J. Scheper, J. B. Wickens, V. J. Wickens, R. Bommarco, D. Kleijn, S. G. Potts, S. P. M. Roberts, H. G. Smith, M. Vilà, A. Vujčić, I. Steffan-Dewenter, Mass-flowering crops dilute pollinator abundance in agricultural landscapes across Europe. *Ecol. Lett.* **19**, 1228–1236 (2016).
50. C. Thies, T. Tschamtkke, Landscape structure and biological control in agroecosystems. *Science* **285**, 893–895 (1999).
51. A. Rusch, R. Chaplin-Kramer, M. M. Gardiner, V. Hawro, J. Holland, D. Landis, C. Thies, T. Tschamtkke, W. W. Weisser, C. Winqvist, M. Woltz, R. Bommarco, Agricultural landscape simplification reduces natural pest control: A quantitative synthesis. *Agric. Ecosyst. Environ.* **221**, 198–204 (2016).
52. J. Chen, Y. Ban, S. Li, China: Open access to Earth land-cover map. *Nature* **514**, 434 (2014).
53. A. Gelman, J. Hill, *Data Analysis Using Regression and Multilevel Hierarchical Models* (Cambridge Univ. Press, 2007).
54. M. van de Pol, J. Wright, A simple method for distinguishing within- versus between-subject effects using mixed models. *Anim. Behav.* **77**, 753–758 (2009).
55. H. Schielzeth, Simple means to improve the interpretability of regression coefficients. *Methods Ecol. Evol.* **1**, 103–113 (2010).
56. B. J. Cardinale, D. S. Srivastava, J. Emmett Duffy, J. P. Wright, A. L. Downing, M. Sankaran, C. Jouseau, Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*. **443**, 989–992 (2006).
57. P.-C. Bürkner, brms: An R package for Bayesian multilevel models using Stan. *J. Stat. Softw.* **80**, 1–27 (2017).
58. D. P. MacKinnon, A. J. Fairchild, M. S. Fritz, Mediation analysis. *Annu. Rev. Psychol.* **58**, 593–614 (2007).
59. D. Lüdtke, sjstats: Statistical functions for regression models. 10.5281/ZENODO.1308979 (2018).
60. B. Carpenter, A. Gelman, M. D. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. Brubaker, J. Guo, P. Li, A. Riddell, Stan: A probabilistic programming language. *J. Stat. Softw.* **76**, 1–37 (2017).
61. J. Kruschke, *Doing Bayesian Data Analysis: A Tutorial with R, JAGS, and Stan* (Academic Press, ed. 2, 2015).
62. R. McElreath, *Statistical Rethinking: A Bayesian Course with Examples in R and Stan* (CRC Press, 2016).
63. A. Vehtari, A. Gelman, J. Gabry, Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Stat. Comput.* **27**, 1413–1432 (2017).
64. A. Vehtari, A. Gelman, J. Gabry, Y. Yao, A. Gelman, loo: Efficient leave-one-out cross-validation and WAIC for Bayesian models. R package version, 2.0.0 2018; <https://cran.r-project.org/package=loo>.
65. U. Samnegård, G. Alins, V. Boreux, J. Bosch, D. García, A. K. Happe, A. M. Klein, M. Miñarro, K. Mody, M. Porcel, A. Rodrigo, L. Roquer-Beni, M. Tasin, P. A. Hambäck, Management trade-offs on ecosystem services in apple orchards across Europe: Direct and indirect effects of organic production. *J. Appl. Ecol.* **56**, 802–811 (2019).
66. M. P. D. Garratt, C. L. Truslove, D. J. Coston, R. L. Evans, E. D. Moss, C. Dodson, N. Jenner, J. C. Biesmeijer, S. G. Potts, Pollination deficits in UK apple orchards. *J. Pollinat. Ecol.* **12**, 9–14 (2014).
67. G. A. de Groot, R. van Kats, M. Reemer, D. van der Sterren, J. C. Biesmeijer, D. Kleijn, “De bijdrage van (wilde) bestuivers aan de opbrengst van appels en blauwe bessen: Kwantificering van ecosystemediensten in Nederland,” Alterra, Wageningen-UR, 2015; <http://edepot.wur.nl/353774>.
68. R. E. Mallinger, C. Gratton, Species richness of wild bees, but not the use of managed honeybees, increases fruit set of a pollinator-dependent crop. *J. Appl. Ecol.* **52**, 323–330 (2015).
69. M. P. D. Garratt, D. J. Coston, C. L. Truslove, M. G. Lappage, C. Polce, R. Dean, J. C. Biesmeijer, S. G. Potts, The identity of crop pollinators helps target conservation for improved ecosystem services. *Biol. Conserv.* **169**, 128–135 (2014).
70. D. d. L. Ramos, M. M. C. Bustamante, F. D. D. S. E. Silva, L. G. Carvalheiro, Crop fertilization affects pollination service provision—Common bean as a case study. *PLOS ONE*. **13**, e0204460 (2018).
71. H. Taki, K. Okabe, S. Makino, Y. Yamaura, M. Sueyoshi, Contribution of small insects to pollination of common buckwheat, a distylous crop. *Ann. Appl. Biol.* **155**, 121–129 (2009).
72. H. Taki, K. Okabe, Y. Yamaura, T. Matsuura, M. Sueyoshi, S. Makino, K. Maeto, Effects of landscape metrics on *Apis* and non-*Apis* pollinators and seed set in common buckwheat. *Basic Appl. Ecol.* **11**, 594–602 (2010).
73. A. Holzschuh, J.-H. Dudenhöffer, T. Tschamtkke, Landscapes with wild bee habitats enhance pollination, fruit set and yield of sweet cherry. *Biol. Conserv.* **153**, 101–107 (2012).
74. V. Boreux, S. Krishnan, K. G. Cheppudira, J. Ghazoul, Impact of forest fragments on bee visits and fruit set in rain-fed and irrigated coffee agro-forests. *Agric. Ecosyst. Environ.* **172**, 42–48 (2013).
75. V. Boreux, P. Vaast, L. P. Madappa, K. G. Cheppudira, C. Garcia, J. Ghazoul, Agroforestry coffee production increased by native shade trees, irrigation, and liming. *Agron. Sustain. Dev.* **36**, 10.1007/s13593-016-0377-7, (2016).



76. A. Classen, M. K. Peters, S. W. Ferger, M. Helbig-Bonitz, J. M. Schmack, G. Maassen, M. Schleuning, E. K. V. Kalko, K. Böhning-Gaese, I. Steffan-Dewenter, Complementary ecosystem services provided by pest predators and pollinators increase quantity and quality of coffee yields. *Proc. Biol. Sci.* **281**, 20133148 (2014).
77. J. Hipólito, D. Boscolo, B. F. Viana, Landscape and crop management strategies to conserve pollination services and increase yields in tropical coffee farms. *Agric. Ecosyst. Environ.* **256**, 218–225 (2018).
78. S. Krishnan, C. G. Kushalappa, R. U. Shaanker, J. Ghazoul, Status of pollinators and their efficiency in coffee fruit set in a fragmented landscape mosaic in South India. *Basic Appl. Ecol.* **13**, 277–285 (2012).
79. S. Krishnan, K. G. Cheppudira, J. Ghazoul, Pollinator services in coffee agroforests of the western Ghats, in *Agroforestry*, J. Dagar, V. Tewari, Eds. (Springer Singapore, 2017), pp. 771–795.
80. M. Nesper, C. Kueffer, S. Krishnan, C. G. Kushalappa, J. Ghazoul, Shade tree diversity enhances coffee production and quality in agroforestry systems in the Western Ghats. *Agric. Ecosyst. Environ.* **247**, 172–181 (2017).
81. S. Cusser, J. L. Neff, S. Jha, Natural land cover drives pollinator abundance and richness, leading to reductions in pollen limitation in cotton agroecosystems. *Agric. Ecosyst. Environ.* **226**, 33–42 (2016).
82. N. P. Chacoff, M. A. Aizen, Edge effects on flower-visiting insects in grapefruit plantations bordering premontane subtropical forest. *J. Appl. Ecol.* **43**, 18–27 (2006).
83. N. P. Chacoff, M. A. Aizen, V. Aschero, Proximity to forest edge does not affect crop production despite pollen limitation. *Proc. R. Soc. B Biol. Sci.* **275**, 907–913 (2008).
84. L. G. Carvalheiro, C. L. Seymour, R. Veldtman, S. W. Nicolson, Pollination services decline with distance from natural habitat even in biodiversity-rich areas. *J. Appl. Ecol.* **47**, 810–820 (2010).
85. L. G. Carvalheiro, C. L. Seymour, S. W. Nicolson, Creating patches of native flowers facilitates crop pollination in large agricultural fields: Mango as a case study. *J. Appl. Ecol.* **49**, 1373–1383 (2012).
86. I. Bartomeus, V. Gagic, R. Bommarco, Pollinators, pests and soil properties interactively shape oilseed rape yield. *Basic Appl. Ecol.* **16**, 737–745 (2015).
87. D. A. Stanley, D. Gunning, J. C. Stout, Pollinators and pollination of oilseed rape crops (*Brassica napus* L.) in Ireland: Ecological and economic incentives for pollinator conservation. *J. Insect Conserv.* **17**, 1181–1189 (2013).
88. D. A. Stanley, J. C. Stout, Pollinator sharing between mass-flowering oilseed rape and co-flowering wild plants: Implications for wild plant pollination. *Plant Ecol.* **215**, 315–325 (2014).
89. Y. Zou, F. J. J. A. Bianchi, F. Jauker, S. Luo, W. van der Werf, Wild pollinators enhance oilseed rape yield in small-holder farming systems in China. *BMC Ecol.* **17**, 6 (2017).
90. H. Connelly, K. Poveda, G. Loeb, Landscape simplification decreases wild bee pollination services to strawberry. *Agric. Ecosyst. Environ.* **211**, 51–56 (2015).
91. H. Grab, K. Poveda, B. Danforth, G. Loeb, Landscape context shifts the balance of costs and benefits from wildflower borders on multiple ecosystem services. *Proc. R. Soc. B Biol. Sci.* **285**, 20181102 (2018).
92. R. I. A. Stewart, G. K. S. Andersson, C. Brönmark, B. K. Klatt, L.-A. Hansson, V. Zülsdorff, H. G. Smith, Ecosystem services across the aquatic-terrestrial boundary: Linking ponds to pollination. *Basic Appl. Ecol.* **18**, 13–20 (2016).
93. B. Ricci, P. Franck, J.-F. Toubon, J.-C. Bouvier, B. Sauphanor, C. Lavigne, The influence of landscape on insect pest dynamics: A case study in southeastern France. *Landsc. Ecol.* **24**, 337–349 (2009).
94. M. Maaloulou, P. Franck, J.-C. Bouvier, J.-F. Toubon, C. Lavigne, Codling moth parasitism is affected by semi-natural habitats and agricultural practices at orchard and landscape levels. *Agric. Ecosyst. Environ.* **169**, 33–42 (2013).
95. B. Caballero-López, R. Bommarco, J. M. Blanco-Moreno, F. X. Sans, J. Pujade-Villar, M. Rundlöf, H. G. Smith, Aphids and their natural enemies are differently affected by habitat features at local and landscape scales. *Biol. Control.* **63**, 222–229 (2012).
96. G. Tamburini, S. De Simone, M. Sigura, F. Boscutti, L. Marini, Conservation tillage mitigates the negative effect of landscape simplification on biological control. *J. Appl. Ecol.* **53**, 233–241 (2016).
97. G. Tamburini, S. De Simone, M. Sigura, F. Boscutti, L. Marini, Soil management shapes ecosystem service provision and trade-offs in agricultural landscapes. *Proc. R. Soc. B Biol. Sci.* **283**, 20161369 (2016).
98. G. Tamburini, I. Pevere, N. Fornasini, S. de Simone, M. Sigura, F. Boscutti, L. Marini, Conservation tillage reduces the negative impact of urbanisation on carabid communities. *Insect Conserv. Divers.* **9**, 438–445 (2016).
99. H. Taki, K. Maeto, K. Okabe, N. Haruyama, Influences of the seminatural and natural matrix surrounding crop fields on aphid presence and aphid predator abundance within a complex landscape. *Agric. Ecosyst. Environ.* **179**, 87–93 (2013).
100. D. K. Letourneau, S. G. Bothwell Allen, J. O. Stireman III, Perennial habitat fragments, parasitoid diversity and parasitism in ephemeral crops. *J. Appl. Ecol.* **49**, 1405–1416 (2012).
101. B. Maas, Y. Clough, T. Tschardtke, Bats and birds increase crop yield in tropical agroforestry landscapes. *Ecol. Lett.* **16**, 1480–1487 (2013).
102. M. E. O. O'Rourke, K. Rienzo-Stack, A. G. Power, A multi-scale, landscape approach to predicting insect populations in agroecosystems. *Ecol. Appl.* **21**, 1782–1791 (2011).
103. M. Jonsson, H. L. Buckley, B. S. Case, S. D. Wratten, R. J. Hale, R. K. Didham, Agricultural intensification drives landscape-context effects on host-parasitoid interactions in agroecosystems. *J. Appl. Ecol.* **49**, 706–714 (2012).
104. M. Kishinevsky, T. Keasar, A. R. Harari, E. Chiel, A comparison of naturally growing vegetation vs. border-planted companion plants for sustaining parasitoids in pomegranate orchards. *Agric. Ecosyst. Environ.* **246**, 117–123 (2017).
105. E. A. Martin, B. Seo, C.-R. Park, B. Reineking, I. Steffan-Dewenter, Scale-dependent effects of landscape composition and configuration on natural enemy diversity, crop herbivory, and yields. *Ecol. Appl.* **26**, 448–462 (2016).
106. K. Poveda, E. Martínez, M. F. Kersch-Becker, M. A. Bonilla, T. Tschardtke, Landscape simplification and altitude affect biodiversity, herbivory and Andean potato yield. *J. Appl. Ecol.* **49**, 513–522 (2012).
107. M. B. Takada, A. Yoshioka, S. Takagi, S. Iwabuchi, I. Washitani, Multiple spatial scale factors affecting mirid bug abundance and damage level in organic rice paddies. *Biol. Control.* **60**, 169–174 (2012).
108. M. G. E. Mitchell, E. M. Bennett, A. Gonzalez, Agricultural landscape structure affects arthropod diversity and arthropod-derived ecosystem services. *Agric. Ecosyst. Environ.* **192**, 144–151 (2014).
109. F. Geiger, J. Bengtsson, F. Berendse, W. W. Weisser, M. Emmerson, M. B. Morales, P. Ceryngier, J. Liira, T. Tschardtke, C. Winqvist, S. Eggers, R. Bommarco, T. Pärt, V. Bretagnolle, M. Plantegenest, L. W. Clement, C. Dennis, C. Palmer, J. J. Oñate, I. Guerrero, V. Hawro, T. Aavik, C. Thies, A. Flohre, S. Hänke, C. Fischer, P. W. Goedhart, P. Inchausti, Persistent negative effects of pesticides on biodiversity and biological control potential on European farmland. *Basic Appl. Ecol.* **11**, 97–105 (2010).
110. M. Plečaš, V. Gagić, M. Janković, O. Petrović-Obradović, N. G. Kavallieratos, Ž. Tomanović, C. Thies, T. Tschardtke, A. Četković, Landscape composition and configuration influence cereal aphid-parasitoid-hyperparasitoid interactions and biological control differentially across years. *Agric. Ecosyst. Environ.* **183**, 1–10 (2014).
111. M. Tschumi, M. Albrecht, M. H. Entling, K. Jacot, High effectiveness of tailored flower strips in reducing pests and crop plant damage. *Proc. R. Soc. Biol. Sci.* **282**, 20151369 (2015).

**Acknowledgments:** We thank R. Carloni for producing icons in Figs. 1 to 4. We also thank the Department of Innovation, Research and University of the Autonomous Province of Bozen/Bolzano for covering the Open Access publication costs. For all further acknowledgements, see the Supplementary Materials. **Funding:** This work was funded by EU-FP7 LIBERATION (311781) and Biodiversa-FACCE ECODEAL (PCIN-2014-048). **Author contributions:** M.D., E.A.M., and I.S.-D. conceived the study. M.D. performed statistical analyses and wrote the manuscript draft. The authors named from E.A.M. to T.T. and I.S.-D. discussed and revised earlier versions of the manuscript. The authors named from G.K.S.A. to Y.Z. are listed alphabetically, as they contributed equally in gathering field data and providing several important corrections to subsequent manuscript drafts. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Additional data related to this paper may be requested from the authors.

Submitted 14 February 2019

Accepted 22 September 2019

Published 16 October 2019

10.1126/sciadv.aax0121

**Citation:** M. Dainese, E. A. Martin, M. A. Aizen, M. Albrecht, I. Bartomeus, R. Bommarco, L. G. Carvalheiro, R. Chaplin-Kramer, V. Gagic, L. A. Garibaldi, J. Ghazoul, H. Grab, M. Jonsson, D. S. Karp, C. M. Kennedy, D. Kleijn, C. Kremen, D. A. Landis, D. K. Letourneau, L. Marini, K. Poveda, R. Rader, H. G. Smith, T. Tschardtke, G. K. S. Andersson, I. Badenhausser, S. Baensch, A. D. M. Bezerra, F. J. J. A. Bianchi, V. Boreux, V. Bretagnolle, B. Caballero-Lopez, P. Cavigliasso, A. Četković, N. P. Chacoff, A. Classen, S. Cusser, F. D. da Silva e Silva, G. A. deGroot, J. H. Dudenhöffer, J. Ekroos, T. Fijen, P. Franck, B. M. Freitas, M. P. D. Garratt, C. Gratton, J. Hipólito, A. Holzschuh, L. Hunt, A. L. Iverson, S. Jha, T. Keasar, T. N. Kim, M. Kishinevsky, B. K. Klatt, A.-M. Klein, K. M. Krewenka, S. Krishnan, A. E. Larsen, C. Lavigne, H. Liere, B. Maas, R. E. Mallinger, E. Martinez Pachon, A. Martínez-Salinas, T. D. Meehan, M. G. E. Mitchell, G. A. R. Molina, M. Nesper, L. Nilsson, M. E. O'Rourke, M. K. Peters, M. Plečaš, S. G. Potts, D. d. L. Ramos, J. A. Rosenheim, M. Rundlöf, A. Rusch, A. Sáez, J. Scheper, M. Schleuning, J. M. Schmack, A. R. Sciligo, C. Seymour, D. A. Stanley, R. Stewart, J. C. Stout, L. Sutter, M. B. Takada, H. Taki, G. Tamburini, M. Tschumi, B. F. Viana, C. Westphal, B. K. Willcox, S. D. Wratten, A. Yoshioka, C. Zaragoza-Trello, W. Zhang, Y. Zou, I. Steffan-Dewenter, A global synthesis reveals biodiversity-mediated benefits for crop production. *Sci. Adv.* **5**, eaax0121 (2019).

## A global synthesis reveals biodiversity-mediated benefits for crop production

Matteo Dainese, Emily A. Martin, Marcelo A. Aizen, Matthias Albrecht, Ignasi Bartomeus, Riccardo Bommarco, Luisa G. Carvalheiro, Rebecca Chaplin-Kramer, Vesna Gagic, Lucas A. Garibaldi, Jaboury Ghazoul, Heather Grab, Mattias Jonsson, Daniel S. Karp, Christina M. Kennedy, David Kleijn, Claire Kremen, Douglas A. Landis, Deborah K. Letourneau, Lorenzo Marini, Katja Poveda, Romina Rader, Henrik G. Smith, Teja Tscharntke, Georg K. S. Andersson, Isabelle Badenhausser, Svenja Baensch, Antonio Diego M. Bezerra, Felix J. J. A. Bianchi, Virginie Boreux, Vincent Bretagnolle, Berta Caballero-Lopez, Pablo Cavigliasso, Aleksandar Cetkovic, Natacha P. Chacoff, Alice Classen, Sarah Cusser, Felipe D. da Silva e Silva, G. Arjen de Groot, Jan H. Dudenhöffer, Johan Ekroos, Thijs Fijen, Pierre Franck, Breno M. Freitas, Michael P. D. Garratt, Claudio Gratton, Juliana Hipólito, Andrea Holzschuh, Lauren Hunt, Aaron L. Iverson, Shalene Jha, Tamar Keasar, Tania N. Kim, Miriam Kishinevsky, Björn K. Klatt, Alexandra-Maria Klein, Kristin M. Krewenka, Smitha Krishnan, Ashley E. Larsen, Claire Lavigne, Heidi Liere, Bea Maas, Rachel E. Mallinger, Eliana Martinez Pachon, Alejandra Martínez-Salinas, Timothy D. Meehan, Matthew G. E. Mitchell, Gonzalo A. R. Molina, Maike Nesper, Lovisa Nilsson, Megan E. O'Rourke, Marcell K. Peters, Milan Plecas, Simon G. Potts, Davi de L. Ramos, Jay A. Rosenheim, Maj Rundlöf, Adrien Rusch, Agustín Sáez, Jeroen Scheper, Matthias Schleuning, Julia M. Schmack, Amber R. Sciligo, Colleen Seymour, Dara A. Stanley, Rebecca Stewart, Jane C. Stout, Louis Sutter, Mayura B. Takada, Hisatomo Taki, Giovanni Tamburini, Matthias Tschumi, Blandina F. Viana, Catrin Westphal, Bryony K. Willcox, Stephen D. Wratten, Akira Yoshioka, Carlos Zaragoza-Trello, Wei Zhang, Yi Zou and Ingolf Steffan-Dewenter

*Sci Adv* 5 (10), eaax0121.  
DOI: 10.1126/sciadv.aax0121

### ARTICLE TOOLS

<http://advances.sciencemag.org/content/5/10/eaax0121>

### SUPPLEMENTARY MATERIALS

<http://advances.sciencemag.org/content/suppl/2019/10/10/5.10.eaax0121.DC1>

### REFERENCES

This article cites 101 articles, 13 of which you can access for free  
<http://advances.sciencemag.org/content/5/10/eaax0121#BIBL>

### PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

---

*Science Advances* (ISSN 2375-2548) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. 2017 © The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. The title *Science Advances* is a registered trademark of AAAS.